

Cedar Key Aquaculture Workshop

Sulfide Concentrations in Sediments and Water: Influence on Hard Clams

Derk Bergquist¹, David Julian², Shirley
Baker³

¹*South Carolina Department of Natural
Resources*




² *University of Florida, Department of Zoology*

³ *University of Florida, Department of Fisheries
and Aquatic Sciences*

Why study sulfide?

- High primary productivity
- Decomposition of organic matter by bacteria uses up oxygen
- Anaerobic decomposition produces hydrogen sulfide
- Sulfide is a metabolic poison
- Known to decrease the growth and survival of many bivalve species




Objectives

- Examine sediment sulfide levels in the Suwannee River Estuary
- Determine the effect of sulfide on hard clam survival




Methods

- Sediment porewater samples collected
 - Inside and outside leases
- Laboratory challenges
 - 2 sizes of clams
 - Hypoxia and sulfide




Sulfide – Field Data







- All lease sites had sediment porewater sulfide
- Clam lease sites had mean sulfide levels of up to 110 $\mu\text{mol/L}$
- Non-lease sites had mean sulfide levels up to 300 $\mu\text{mol/L}$

Sulfide – Field Data



Sulfide levels greatest (up to 400 $\mu\text{mol/L}$) in August and September when temperatures high

Laboratory Challenges - Sulfide levels in sediments can affect survival of clams



Sulfide - Conclusions

- Sulfide occurs at lease sites
- Sulfide is greatest in August & September
- Seed clams vulnerable to sulfide toxicity in the lab
- More Questions
 - Does sulfide cause losses in the field?
 - Is sulfide present in the incurrent water?
 - Are larger clams more or less vulnerable?

